

# Hausdorff dimension and filling factor

Wellington da Cruz\*

*Departamento de Física,*

*Universidade Estadual de Londrina, Caixa Postal 6001,*

*Cep 86051-970 Londrina, PR, Brazil*

(February 1, 2008)

## Abstract

We propose a *new hierarchy scheme* for the *filling factor*, a parameter which characterizes the occurrence of the Fractional Quantum Hall Effect ( FQHE ). We consider the Hausdorff dimension,  $h$ , as a parameter for classifying fractional spin particles, such that, it is written in terms of the statistics of the collective excitations. The number  $h$  classifies these excitations with different statistics in terms of its homotopy class.

PACS numbers: 12.90+b

Keywords: Hausdorff dimension; Fractional spin particles; Filling factor

Typeset using REVTeX

---

\*E-mail: wdacruz@fisica.uel.br

In a series of papers [1], we have obtained a set of results with respect to fractional spin particles. Now we make the connection with the filling factor, a parameter that appears in the context of the FQHE and for it we have experimental values [2]. That connection can be made once the anyonic model has been considered to explain this phenomenon [3].

The FQHE is associated with a planar charged system in a perpendicular magnetic field such that a new type of correlated ground state occurs. The Hall resistance develops plateaus at quantized values in the vicinity of the filling factor or statistics,  $\nu$ , which is related to the fraction of electrons that forms collective excitations as quasiholes or quasiparticles. Excitations above the Laughlin ground state are characterized, therefore, by  $\nu$  and so we propose a *new hierarchy scheme* for the FQHE which gives us the possibility of *predicting* for which values of  $\nu$  FQHE can be observed.

Our scheme is based on the intervals of definition of spin,  $s$ , for fractional spin particles which are related to the Hausdorff dimension,  $h$ . We verify that for some experimental values of  $\nu$  for which the FQHE was observed the Hausdorff dimension is a rational number with an odd denominator ( like the filling factor ). Thus, bearing in mind the condition,  $1 < h < 2$ , we can determine for which values of  $h$ , the statistics give numbers in the intervals of definition, as follows:

$$\begin{aligned}
h_1 &= 2 - \nu, & 0 < \nu < 1; & & h_2 &= \nu, & 1 < \nu < 2; \\
h_3 &= 4 - \nu, & 2 < \nu < 3; & & h_4 &= \nu - 2, & 3 < \nu < 4; \\
h_5 &= 6 - \nu, & 4 < \nu < 5; & & h_6 &= \nu - 4, & 5 < \nu < 6; \\
h_7 &= 8 - \nu, & 6 < \nu < 7; & & h_8 &= \nu - 6, & 7 < \nu < 8; \\
h_9 &= 10 - \nu, & 8 < \nu < 9; & & h_{10} &= \nu - 8, & 9 < \nu < 10; \\
&& & & & & etc.
\end{aligned} \tag{1}$$

In these formulas,  $h_i$ , represents the Hausdorff dimension of the collective excitations which are characterized by the statistics  $\nu$ . This *hierarchy scheme* confirms our observation that *when the particles are interacting, in the presence of Chern-Simons field, the Hausdorff dimension changes because the statistics of the collective excitations change* [1]. Now, we give for some experimental values of  $\nu$  the respective values of  $h$ , that is,

$$(h, 0 < \nu < 1) : \tag{2}$$

$$\begin{aligned}
&\left(\frac{9}{5}, \frac{1}{5}\right), \left(\frac{12}{7}, \frac{2}{7}\right), \left(\frac{5}{3}, \frac{1}{3}\right), \left(\frac{8}{5}, \frac{2}{5}\right), \\
&\left(\frac{11}{7}, \frac{3}{7}\right), \left(\frac{14}{9}, \frac{4}{9}\right), \left(\frac{13}{9}, \frac{5}{9}\right), \left(\frac{10}{7}, \frac{4}{7}\right), \\
&\left(\frac{7}{5}, \frac{3}{5}\right), \left(\frac{4}{3}, \frac{2}{3}\right), \left(\frac{6}{5}, \frac{4}{5}\right);
\end{aligned}$$

$$(h, 1 < \nu < 2) : \tag{3}$$

$$\begin{aligned}
&\left(\frac{5}{3}, \frac{5}{3}\right), \left(\frac{13}{7}, \frac{13}{7}\right), \left(\frac{13}{9}, \frac{13}{9}\right), \\
&\left(\frac{10}{7}, \frac{10}{7}\right), \left(\frac{7}{5}, \frac{7}{5}\right), \left(\frac{4}{3}, \frac{4}{3}\right);
\end{aligned}$$

$$(h, 2 < \nu < 3) : \tag{4}$$

$$\left(\frac{4}{3}, \frac{8}{3}\right), \left(\frac{5}{3}, \frac{7}{3}\right).$$

We can see another interesting point from these pairs of numbers: Some collective excitations with different spins have the same value of  $h$ , that is, the nature of the occurrence of FQHE for that values of  $\nu$  can be classified in terms of  $h$ , so we can say that this number classifies *the collective excitations in terms of its homotopy class* [3]. In this way, the Laughlin wavefunctions [4] can be understood as a mapping between homotopy classes of the collective excitations.

## ACKNOWLEDGMENTS

I would like to thank Steven F. Durrant for reading the manuscript.

## REFERENCES

- [1] W. da Cruz, preprint/UEL-DF/W-01/97 (submitted to Phys. Rev. D. ); W. da Cruz, preprint/UEL-DF/W-02/97; W. da Cruz, preprint/UEL-DF/W-03/97 ( hep-th/9802135 ).
- [2] A. M. Chang in *The Quantum Hall Effect*, ed. by R. E. Prange and S. M. Girvin, ( Springer Verlag, 1987) and references therein.
- [3] A. Lerda, *Anyons*, Lectures Notes in Physics, ( Springer Verlag, 1992); S. Forte, Rev. Mod. Phys. **64**, 193 (1992) and references therein.
- [4] R. B. Laughlin, Phys. Rev. Lett.**50** 1395 (1983); *ibid*, Phys. Rev. **B23** 3383 (1983); B. I. Halperin, Phys. Rev. Lett. **52** 1583 (1984); F. D. M. Haldane, Phys. Rev. Lett. **51** 605 (1983); R. Tao and D. J. Thouless, Phys. Rev. **B28** 1142 ( 1983); P. W. Anderson, Phys. Rev. **B28** 2264 (1983); J. K. Jain, Adv. Phys. **41**, 105 (1992) and references therein; J. E. MacDonald and F. D. M. Haldane, Phys. Rev. **B55**, 7818 (1997); R. Mélin and B. Douçot, Phys. Rev. **B55** 13135 (1997) and references therein.